

HIGHLY RECOMMENDED

New &
Updated

 **OSWAAL BOOKS®**
LEARNING MADE SIMPLE

Handbook MATHEMATICS

Useful For

Classes 11 & 12

All Leading Competitive Exams

#1 Mathematics Toolkit

1

**Get Concept Clarity
& Revision**

with Important
Formulae & Derivations

2

**Fill Learning
Gaps**

with 300+ Concept
Videos

3

**Get Valuable
Concept Insights**

with Appendix,
Smart Mind Maps
& Mnemonics

4

**Free Online
Assessment**

with
OSWAAL 360°



2nd EDITION

YEAR 2023-24



ISBN

“978-93-89510-20-1”



**SYLLABUS
COVERED**

CBSE, CISCE,
JEE Mains and Advanced



**COPYRIGHT
RESERVED
BY THE PUBLISHERS**

All rights reserved. No part of this book may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, without written permission from the publishers. The author and publisher will gladly receive information enabling them to rectify any error or omission in subsequent editions.



PUBLISHED BY



**OSWAAL BOOKS &
LEARNING PVT. LTD.**



1/11, Sahitya Kunj, M.G. Road,
Agra - 282002, (UP) India



1/1, Cambourne Business Centre
Cambridge, Cambridgeshire
CB 236DP, United kingdom



0562-2857671



contact@oswaalbooks.com



www.OswaalBooks.com

DISCLAIMER

This book is published by Oswaal Books and Learning Pvt Ltd ("Publisher") and is intended solely for educational use, to enable students to practice for examinations/tests and reference. The contents of this book primarily comprise a collection of questions that have been sourced from previous examination papers. Any practice questions and/or notes included by the Publisher are formulated by placing reliance on previous question papers and are in keeping with the format/pattern/guidelines applicable to such papers.

The Publisher expressly disclaims any liability for the use of, or references to, any terms or terminology in the book, which may not be considered appropriate or may be considered offensive, in light of societal changes. Further, the contents of this book, including references to any persons, corporations, brands, political parties, incidents, historical events and/or terminology within the book, if any, are not intended to be offensive, and/or to hurt, insult or defame any person (whether living or dead), entity, gender, caste, religion, race, etc. and any interpretation to this effect is unintended and purely incidental. While we try to keep our publications as updated and accurate as possible, human error may creep in. We expressly disclaim liability for errors and/or omissions in the content, if any, and further disclaim any liability for any loss or damages in connection with the use of the book and reference to its contents".

Kindle Edition

Order Form

ONE FOR ALL OLYMPIADS (Multicolour)

Classes 1 to 5

Mathematics, English, Reasoning, General Knowledge, Science & Cyber

ONE FOR ALL WORKBOOKS CONCEPT-WISE (Multicolour)

Classes 1 to 5

Mathematics, English, General Knowledge, Science

NCERT WORKBOOKS (Multicolour / B W)

Classes 1 & 2

English, Hindi & Mathematics

Classes 3 to 5

English, Hindi, Mathematics & EVS

ONE FOR ALL OLYMPIADS

Classes 6 to 8

Mathematics, English, Reasoning, General Knowledge, Science & Cyber

NCERT/CBSE PULL-OUT WORKSHEETS

Classes 6 to 8

Mathematics, English, Science, Social Science, Hindi & Sanskrit

Classes 9 & 10

English Lang. & Literature, Science, Social Science & Mathematics

CBSE QUESTION BANKS (CHAPTER-WISE & TOPIC-WISE SOLVED PAPERS)

Classes 6 to 8

Mathematics, English, Science, Social Science, Hindi & Sanskrit

CBSE QUESTION BANKS (CHAPTER-WISE & TOPIC-WISE SOLVED PAPERS)

Classes 9 & 10

Hindi-A, Hindi-B, English Lang. & Lit. (With NCERT Workbook), Science, Mathematics, Social Science, Computer Applications, Sanskrit & English Communicative (With CBSE Textbook),

Classes 11 & 12

Hindi Core, English Core, Mathematics, Chemistry, Applied Mathematics, Biology, Physics, Accountancy, Economics, Business Studies, Geography, History, Political Science, Computer Science, Entrepreneurship, Informatics Practices, Physical Education, Sociology (For Class 12th Only), Psychology (For Class 12th Only)

NCERT NCERT EXEMPLAR PROBLEMS-SOLUTIONS (Hindi & English Medium)

Classes 9 & 10

Science, Mathematics, गणित, विज्ञान

NCERT NCERT EXEMPLAR PROBLEMS-SOLUTIONS (Hindi & English Medium)

Classes 11 & 12

Physics, Chemistry, Mathematics, Biology, गणित, भौतिक विज्ञान, रसायन विज्ञान, जीव विज्ञान

NCERT TEXTBOOK+EXEMPLAR

Classes 11 & 12

Physics, Chemistry, Mathematics & Biology

CBSE 6 YEARS SOLVED PAPERS

Class 10

English Lang. & Lit, Science, Social Science, Hindi-B, Mathematics Basic, Mathematics Standard, Hindi-A, Computer Applications & Sanskrit

Class 12

Science Stream- (Chemistry, Physics, Mathematics, English Core & Biology), Commerce Stream- Economics, (Accountancy, Business Studies, Mathematics & English Core), Humanities Stream- (Geography, History, Political Science, English Core & Hindi Core)

CBSE RMT FLASH CARD

Class 10

Mathematics, English Lang. & Lit., Social Science & Science

CBSE ONE FOR ALL

Classes 9 & 10

Mathematics, English Lang. & Lit., Social Science, Science, Hindi-A, Hindi-B, Mathematics (Stand.) & Mathematics Basic

Class 12

English Core, Physics, Chemistry, Mathematics, Biology, Accountancy, Business Studies & Economics

CBSE LMP-Last Minute Preparation System

Class 10

Combined (English Lang. & Lit., Science, Mathematics Standard & Social Science)

Class 12

Science (Physics, Chemistry, Mathematics, Biology & English Core)
Commerce (Accountancy, Business Studies, Economics, Mathematics & English Core)
Humanities (History, Geography, Political Science & English Core)

CBSE SAMPLE QUESTION PAPERS

Classes 9 & 10

Hindi-A, Hindi-B, English Lang. & Lit., Computer Appl., English Communicative, Mathematics, Science, Social Science, Sanskrit & Samajik Vigyan (For Class 10th Only)

Classes 11 & 12

Hindi Core, English Core, Mathematics, Applied Mathematics, Physics, Chemistry, Biology, Accountancy, Economics, Business Studies, Geography, History, Political Science, Computer Science, Informatics Practices, Physical Education, Entrepreneurship (For Class 12th Only), Psychology (For Class 12th Only) & Sociology (For Class 12th Only)

Exam Preparation Books for Class 1 to 12 : CBSE, CISCE Boards & Karnataka Board | OLYMPIADS | JEE | NEET | CUET | NDA | CDS | CAT | GATE | UPSC | UGC NET | NRA CET & more

We are available in all leading e-commerce platforms and all good bookstores.



Exclusive School Books Suppliers

VIJAYAWADA	ANDHRA PRADESH Sri Vikas Book Centre, 9848571114, 9440715700, Sri Kanka Durga Book Stall, 9603491009
	ASSAM Dutta Book Stall, 8729948473
WEST KAMENG	JHARKHAND Bharti International, 06432359136, 9431132051
DEOGHAR	KARNATAKA Palasandra Promoters, 9448214377, (0816) 2273220
TUMKUR BANGLORE	Sree byraveshawara book Depot, 9880561081,
INDORE	MADHYA PRADESH Adithi Distributor, 9229171880
KOLHAPUR	MAHARASHTRA Ashish Book Depot, 7798420420

PUNE	Pravin Sales, 9890683475, Sai Shubham, 9975687687
DHULE	Navjeevan Book Stall, 9422296434
JALNA	Anil Paper Mart, 9422722522, (02482) 230733
AGRA	UTTAR PRADESH Maheshwari Pustak Bhandar, 9760602503
	TAMIL NADU Bookmark-IT, 7305151653
CHENNAI	TELANGANA
HYDERABAD	Sri Balaji Book Depot , 9676996199, (040) 27613300, Shri Sharada Store & Stationers, 9000400069
KOLKATA	WEST BENGAL
WEST KEVVERY	United Book House, 9831344622 Schoolwale & Company, 9731715655

Our Distributors

VISAKHAPATHAM	ANDHRA PRADESH JBD Educational, 9246632691, 9246633693
	ANDAMAN & NICOBAR ISLAND Krishna Book Centre, 9474205570
PORTBLAIR	ASSAM National Library, 9435031757
DIBRUGARH GOALPARA GUWAHATI	Sanjay Book Centre, 9706963878 Book Emporium, 9675972993, 6000763186, Ashok Publication, 7896141127, Kayaan Enterprises, (0361) 2630443, Central Book Agency, 99547107165, Orchid Book house, 9864624209
PATNA	BIHAR Bokaro Student Friends, (0612) 2300600, 2618400, Gyan Ganga Ltd., 9304826651 Nova Publisher & Distributors, (0612) 2666404, Shri Durga Pustak Mandir, 9334477386, Sharda Pustak Bhandar, 9334259293, Vikas Book Depot, 9504780402
MADHUBANI PURNEA	Krishna Book Agency, 9801019292 Chaurasia book centre, 7004456102
BHILAI RAIPUR RAIGARH	CHATTISGARH Anil Book Depot, 9425234260, 8305247406 Shri Ramdev Traders, 9981761797 Sindhu Book Depot, 9425275763, 9981935763
DELHI	DELHI Mittal Books, (011) 23288887, 9899037390, Shyam Brother, 9313619935, Bokaro.Student Friends Pvt Ltd., 7004074900, R. D. Chawla & Sons, 9899445522
GOA	GOA Golden Heart Emporium, (0832) 2725208, 9370273479
AHMEDABAD	GUJARAT Patel Book, 9898184248, 9824386112, 9825900335, Shalibhadra Stationers, 9904477855, Zaveri Agency, 9979897312, 9979890330 Ajay Book Stall (Vidya Nagan), 9723821016
ANAND	Goutam Book Sellers, 9081790813
VAPI	Mahavir Stationers, 9429474177
VALSAD	College Store, 8141913750, 9825099121
NAVSARI	Kazi Brothers, 7984880995, Shopping Point, 0261-2230097
SURAT	
ROHTAK	HARYANA Manish Traders, 9812556687, Swami Kitab Ghar, 9255121948
BALLABGARH FARIDABAD	Kashi Ram Kishan Ial, 9289504004, 8920567245 Agarwal Book House, 9810657456, Ayush Prakashan, 9871529692
BOKARO DHANBAD RANCHI	JAMMU Book Place, 8082160733
DUMKA	JHARKHAND Bokaro Student Friends, (0654) 2233094, 7360021503 Bokaro Student Friends, (0326) 3590527
DEOGHAR	Bokaro Student Friends, 9234628152, Crown Book Distributor & Publishers, (0651) 2213735, 9431173904, Jayshree Book House, 9431350775, Gyan Ganga Ltd., 9117889900, Pustak Mandir, 9431115138, Vidyantathi Pustak Bhandar, 9431310228
HUBLI BANGLORE BELLARY KARWAR	Navyug Sahitya Mandir, 9334375509, 9431367981 KARNATAKA Renuka Book Distributor, (0836) 2244124 Hema Book world, 8040105110, Krishna book house, 9739847334 Chaitanya Exhibition, 9886393971 Akshita Marketing, 9945891555
ERNAKULAM	KERALA Asad Book Centre, (0484) 2370431, 9447314548, Academic Book House, (0484) 2376613, Surya Book Centre, 9847238314, H & C Store, 9864196344, Surya Book House, 9847124217, 9847238314
KOTTAYAM	Book Centre, (0481) 2566992
TRIVANDRUM	Academic Book House, (0471) 2333349, 9447063349, T.B.S. Publisher Distributors, (0471) 2570504 T.B.S. Publisher Distributors, 755982223
KOZHIKODE CALICUT	Aman Book Stall, (0495) 2721282,
KOLLAM PERUMBAAVOOR	Vasantha Book Stall, 9895072023 Adithya book house, 8606005600
GWALIOR INDORE	MADHYA PRADESH Agarwal Book Depot, 9425116210 Arun Prakashan, 9424890785, Bhaiya Industries, 9893326853, Bharti Publishers and Distributors, 7000989901, Sushil Prakashan,(0731) 2503333, 2535892, 9425322330, Bhaiya Store, 9425318103, Vinay Pustak Sadan, 8962362667
JABALPUR	
SAGAR	New Radhika Book Palace, 9425411533-66, Vidya Sagar Stationery Mart, 9425451565
AMRAVATI PUNE	MAHARASHTRA Delta Pen House, 9890350073 Natraj Book Depot, (020) 24485054, 9890054092, Vikas Book House, 9921331187, Sai Shubham, 9975687687, 9890043496, Pravin Sales, 9890683475, New Saraswati Granth Bhandar, 9422323859, Akshar Books & Stationery, 7385089789, Vandhaman Educational, 9860574354, The Wisdom Store, 9822588775, 02025433344 Hira Stationers, 9423755472
AHMED NAGAR	

MUMBAI	Vidyarathi Sales Agencies, 9819776110, New Student Agencies, 7045065799, Shivam Books & Stationery, (022) 28230897, 9892935799
LATUR KOLHAPUR	Yash Book House, 9637936999 Granth the Book World, 9922295522, New India book House, 9623123458, 7448123456
NASHIK NAGPUR	Laxmi Pustakalay and Stationers, (0712) 2727354 Vijay Book Depot, 9860122094, (0712) 2534217, Renuka Book distributor, 9765406133, Novelty Book Depot, 9657690220
DHULE YAVATMAL	Navjeevan Book Stall, 7020525561 Shri Ganesh Pustkalaya, 9423131275
CUTTACK BHUBANESHWAR	ODISHA A. K. Mishra Agencies, 9437025991, 9437081319 M/s Pragnya, 8847888616, 9437943777, Padmalaya, 9437026922, Bidyashree, 9937010700
BARIPADA KEONJHAR BRAHMAPUR	Trimurti Book World, 9437034735 Students corner, 7008435418 Sibani Book Store, 8763340822
BHATINDA PATIALA	PUNJAB Agarwal Book Centre, 9417816439 Adarsh Entreprises, 9814347613
JALANDHAR	Cheap Book Store, 9872223458, 9878258592, City Book Shop, 9417440753, Gaurav Book World, 9478821183, Subhash Book Depot, 9876453625
LUDHIANA	Gupta Brother, 988200206, Amit Book, 9815807871
CHANDIGARH	Mahindra Book Depot, 9814920226
KOTA	RAJASTHAN Perfect Stationers & Gen. Shoppe, 9829863904, Vardhman Book Depot, 9571365020, Bhandari Stationers, 9001094271 Nakoda Book Depot, (01482) 243653, 9214983594
BHILWARA	
JAIPUR	J K Enterprises, 9460983939, 9414782130, Ravi Enterprises, 9829060694, Saraswati Book House, (0141) 2610823, 9829811155
AGARTALA	TRIPURA Book Corner, 8794894165, 8984657146
COIMBATORE CHENNAI	TAMIL NADU Majestic Book House, (0422) 2384333 Arraba Book Traders, (044) 25387868, 9841459105, M.R. Book Store (044) 25364596, Indian Book House, (044) 24327784, 9094129595, Kalaimagal Store, (044) 5544072, 9940619404, Vijaya Stores, 9381037417, Bookmark It-Books & Stat. Store, 7305151653, M.K. Store, 9840030099, Tiger Books Pvt. Ltd., 9710447000
PUDUCHERRY	Sri Saraswathi Book Stall, (04132) 222283, 9092626287
SALEM TRICHY THENI	Pattu book centre, 9894816280 Rasi Publication, 9894816280 Maya Book Centre, 9443929274
HYDERABAD	TELANGANA Sri Balaji Book Depot, (040) 27613300, 9866355473, Central Book Shop, 9121152053, Shah Book House, 9849564564
SECUNDERABAD	Shraddha Books & Stationery, 9849451558
AGRA	UTTAR PRADESH Sunjay Publications, 8126699922, Sparsh Book Agency, 9412257817, Manav Book world, 9258429663, Om Pustak Mandir, (0562) 2464014, 9319117771, Panchsheel Books, 9412257961, 9412257962
ALLAHABAD AZAMGARH ALIGARH	Mehrotra Book Agency, (0532) 2266865, 9415636890 Sasta Sahitya Sadan, 9450029674, Shree Pustak Vatika, 6392555866
BAREILLY BANDA	K.B.C.L. Agarwal, 9897124960
ETAWAH GORAKHPUR JHANSI	Deepak Book store, 9837027416, Bunty Stationers, 9412048717, Swastik Traders, 9415437971
KANPUR	Radhika Competition, 8923681800 Central Book House, 9935454590 Bhanu Book Depot, 9415031340
LUCKNOW	Radha News Agency, 8957247427, Raj Book Dist., 9235616506, H K Book Dist., 9506033137, Bhatia Education & Book Agencies, 9140286011
MEERUT	Vyapar Sadan, 7607102462, Om Book Depot, 7705871398, Vidyarathi kendra Publisher & Distribu- tor, 9554967415, Azad Book Depot Pvt.Ltd., 7317000250, Book Sadan, 9839487327, Universal Books, 8881093333, 7652043110
MATHURA NOIDA	Ideal Book Depot, (0121) 4059252, 9837066307
RENUKOOT VARANASI	Goyal Book Center, 9458766132, M/S Mukesh Pustak Bhandar, 8923338205, Prozo (Global Edu4 Share Pvt. Ltd), 9318395520, Goyal Books Overseas, (0120) 4655555, 9873387003
DEHRADUN	Om Stationers, 9838186991 Bokaro Student Friends, (0542) 2401250, 8299344607, Bookman & Company, 9935194495, Gupta Books, 9918155500
KOLKATA	UTTARAKHAND Brother Pustak Bhandar, 9897833882
COOCH BEHAR	WEST BENGAL
SILIGURI	Oriental Publishers & Distributor (033) 40628367, Katha 'O' Kahini, (033) 22196313, 22419071, Saha Book House, (033), 22193671, 9333416484, United Book House, 9831344622, Bijay Pustak Bhandar, 8961260603, Krishna Book House, 8420431083
	S.B. Book Distributor, Cooch behar, 9002670771
	Modern Book Agency, 8145578772

PREFACE

There are numerous competitive exams conducted in India every year. In this age of cut throat competition, every student aspires to crack the ones he/she appears for in order to begin the journey towards his desired career. Students preparing for various competitive exams strictly follow the prescribed syllabus which has details about the official blueprint as well as typology of questions asked year after year. Along with this, what remains at the core of their preparation is the clarity of concepts.

Oswaal Books is happy to announce the launch of Oswaal Handbooks for Physics, Chemistry, Biology & Mathematics which will supplement the need for concept clarity at every step of study. With the motto of Learning Made Simple, Oswaal Books by the way of these handbooks, aim at providing the most relevant content to students to make studying even the most difficult concepts, interactive and fun.

Oswaal Handbooks cover each and every concept which may be required for the preparation of XI – XII (CBSE, CISCE, IGCSE and all State – Boards), NEET, JEE (Mains and Advanced) and all State and Central Governments Services Exams.

The Handbooks aim to be a ‘Ready Reckoner’ for the students which will help them at every step with every query that they might have.

Some key highlights of Oswaal Handbooks are:

- **Get Concept Clarity & Revision** - with Important Formulae & Derivations.
- **Fill Learning Gaps** - with 300+ Concept Videos.
- **Get Valuable Concept Insights** - with Appendix, Smart Mind Maps & Mnemonics.
- **Free Online Assessment** - with oswaal 360°

We have tried to make our books 100% error-free. Yet, we shall be grateful to the users who point out any errors or make suggestions for the betterment of this book.

We hope that OSWAAL HANDBOOKS will help you at every step as you move closer to your educational goal. We wish you all great success ahead!

All the Best!!
TEAM OSWAAL



CONTENTS

■ Syllabus	7 - 8
■ Mnemonics	9 - 16
<hr/>	
1. Sets and Representations	1 - 6
2. Relations and Functions	7 - 18
3. Trigonometry	19 - 32
4. Complex Numbers	33 - 39
5. Quadratic Equations	40 - 45
6. Linear Inequalities	46 - 51
7. Principle of Mathematical Induction	52 - 56
8. Sequences and Series	57 - 63
9. Permutations and Combinations	64 - 69
10. Binomial Theorem	70 - 76
11. Straight Lines	77 - 82
12. Conic Sections	83 - 95
13. Three Dimensional Geometry	96 - 106
14. Vector Algebra	107 - 113
15. Limits and Derivatives	114 - 120
16. Continuity, Differentiability and Application of Derivatives	121 - 127
17. Determinants	128 - 135
18. Matrices	136 - 143
19. Integral and its Applications	144 - 153
20. Differential Equations	154 - 160
21. Mathematical Reasoning	161 - 165
22. Statistics	166 - 178
23. Commercial Mathematics	179 - 182
24. Probability	183 - 192
➤ Appendix - A	193 - 194
➤ Appendix - B	195 - 195
➤ Appendix - C	195 - 221
➤ Appendix - D	221 - 222
➤ Appendix - E	223 - 223

□□



SYLLABUS

- Sets and Representations** : Sets and their representations, Empty set, Finite and Infinite sets, Equal sets, Subsets of a set of real numbers especially intervals (with notations), Power set, Universal set, Venn diagrams, Union and Intersection of sets, Difference of sets, Complement of a set, Properties of Complement set, Algebraic properties of Union, Intersection and Complement of sets.
- Relations and Functions** : Ordered pairs, Cartesian product of sets, Number of elements in the Cartesian product of two finite sets, Cartesian product of the set of real with itself (upto $\mathbb{R} \times \mathbb{R} \times \mathbb{R}$), Definition of relation, pictorial diagrams, domain, co-domain and range of a relation, types of relation, binary operation, Types of relations: reflexive, symmetric, transitive and equivalence relations. One to one and onto functions, composite functions, inverse of a function. Binary operations. Function as a special type of relation, Pictorial representation of a function, domain, co-domain and range of a function, real valued functions, domain and range of these functions, constant, identity, polynomial, rational, modulus, signum, exponential, logarithmic and greatest integer functions, with their graphs, composition of function, sum, difference, product and quotients of functions.
- Trigonometry**: Trigonometric Functions
 - Positive and negative angles
 - Measuring angles in radians and in degrees and conversion from one measure to another
 - Definition of trigonometric functions with the help of unit circle
 - Truth of the identity $\sin^2 x + \cos^2 x = 1$, for all x
 - Signs of trigonometric functions
 - Domain and range of trigonometric functions and their graphs
 - Expressing $\sin(x \pm y)$ and $\cos(x \pm y)$ in terms of $\sin x$, $\sin y$, $\cos x$ & $\cos y$ and their simple applications
 - Deducing identities like the following :

$$\tan(x \pm y) = \frac{\tan x \pm \tan y}{1 \mp \tan x \cdot \tan y}, \cot(x \pm y) = \frac{\cot x \cdot \cot y \mp 1}{\cot y \pm \cot x}$$
 - $\sin \alpha \pm \sin \beta = 2 \sin \frac{1}{2}(\alpha \pm \beta) \cdot \cos \frac{1}{2}(\alpha \mp \beta)$
 - $\cos \alpha + \cos \beta = 2 \cos \frac{1}{2}(\alpha + \beta) \cdot \cos \frac{1}{2}(\alpha - \beta)$
 - $\cos \alpha - \cos \beta = -2 \sin \frac{1}{2}(\alpha + \beta) \cdot \sin \frac{1}{2}(\alpha - \beta)$
 - Double angle, triple angle, half angle and one third angle formula as special cases.
 - Identities related to $\sin 2x$, $\cos 2x$, $\tan 2x$, $\sin 3x$, $\cos 3x$ and $\tan 3x$
 - Angle and Arc lengths
 - Graphs of simple trigonometric functions for all trigonometric ratios
 - Addition and subtraction formula : $\sin(A \pm B)$; $\cos(A \pm B)$; $\tan(A \pm B)$; $\tan(A + B + C)$ etc
 - Sum and differences as products
 - Product to sum or difference
 - Trigonometric equations: General solution of trigonometric equations of the type $\sin y = \sin x$, $\cos y = \cos x$ and $\tan y = \tan x$
 - Properties of triangle in terms of
 - Sine formula: $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$
 - Cosine formula: $\cos A = \frac{b^2 + c^2 - a^2}{2bc}$, etc.
- Area of triangle $= \frac{1}{2} bc \sin A$ etc.
- Inverse Trigonometric Functions : Definition, range, domain, principal value branch. Graphs of inverse trigonometric functions. Elementary properties of inverse trigonometric functions.
- Complex Numbers**: Need for complex numbers, especially $\sqrt{-1}$, to be motivated by inability to solve some of the quadratic equations, algebraic properties of complex numbers, conjugate complex numbers, Argand plane and polar representation of complex numbers, modulus and argument, triangle inequality, loci on the Argand Diagram, product and quotient for complex numbers in modulus-argument form, De Moivre's theorem, square root of complex number, cube root of unity.
- Quadratic Equations** : Statement of Fundamental Theorem of Algebra, Solution of quadratic equations (with real coefficients), Equations reducible to quadratic form, Nature of roots – Product and sum of roots, Framing of quadratic, cubic, biquadratic equation with given roots, Conditions for common roots in quadratic equations, Properties of quadratic equation, Understanding the fact that a quadratic expression (when plotted on a graph) is a parabola, Minimum and Maximum value of quadratic equations, Sign when the roots are real and when they are complex, Interval of roots, Algebraic interpretation of Rolle's theorem, Descartes' rule of sign.
- Linear Inequalities** : Linear inequalities, Algebraic solutions of linear inequalities in one variable and their representation on the number line. Graphical solution of linear inequalities in two variables. Graphical method of finding a solution of system of linear inequalities in two variables.
- Principle of Mathematical Induction** : Process of the proof by induction, motivating the application of the method by looking at natural numbers as the least inductive subset of real numbers. The principle of mathematical induction and simple applications.
- Sequences and Series**: Sequence and series, Arithmetic Progression (AP), Arithmetic Mean (AM), Geometric Progression (GP), General Term of a GP, Sum of n terms of a GP, infinite GP and its sum, Geometric Mean (GM), Harmonic Progression (HP), n th term of Harmonic Progression, Harmonic Mean, Relation in AM, GM and HM
 - Arithmetic - Geometric Progression (AGP), Sum of Arithmetic-Geometric Series
 - Formulae for the following special sums.

$$\sum_{k=1}^{k=n} k, \sum_{k=1}^{k=n} k^2 \text{ and } \sum_{k=1}^{k=n} k^3$$
- Permutations and Combinations** : Fundamental Principle of Counting, Factorial and its properties, Permutation and its properties, Circular Permutation, Combination and its properties, Exponent of Prime p in $n!$, Division into Groups, use of permutation and combination in geometry, use of permutation and combination in evaluating prime factor, use of permutation and combination in Integral solution, use of permutation combination in Sum of Digits, Derangements theorem.
- Binomial Theorem**: History, statement and proof of the binomial theorem for positive integral indices. Properties of binomial coefficients. Pascal's Triangle, General and middle term in binomial expansion, Binomial Theorem for Any Index, Simple Applications.
- Straight Lines**: Brief recall of two dimensional geometry from earlier classes, shifting of origin, slope of a line and

.....Contd Syllabus

angle between two lines, Various forms of equations of a line, parallel to axis, point-slope form, slope-intercept form, two-point form and normal form, general equation of a line, equation of family of lines passing through the point of intersection of two lines, distance of a point from a line, angle between two lines, equation of lines bisecting the angle between two lines, condition for concurrence of three lines, coordinates of centroid, orthocenter and circumcenter of a triangle

12. **Conic Sections:** Sections of a cone: circles, ellipse, parabola, hyperbola, a point a straight line and a pair of intersecting lines as a degenerated case of a conic section. Standard equations and simple properties of parabola, ellipse and hyperbola, Standard equation of a circle.
13. **Three Dimensional Geometry:** Coordinate axes and coordinate planes in three dimensions. Coordinates of a point. Distance between two points and section formula, direction cosines and direction ratios of a line joining two points, cartesian equation and vector equation of a line, coplanar and skew lines, shortest distance between two lines. Cartesian and vector equation of a plane. Intersection of a line and a plane in different form, angle between (i) two lines, (ii) two planes, (iii) a line and a plane, distance of a point from a plane.
14. **Vector Algebra:** Derivative introduced as rate of change both as that of distance function and geometrically, Intuitive idea of limit, Fundamental theorems on limits, Limits of polynomials and rational functions trigonometric, exponential and logarithmic functions, Definition of derivative relate it to slope of tangent of the curve, Derivative of sum, difference, product and quotient of functions, Derivatives of polynomial and trigonometric functions, Derivative of composite functions, Parametric functions, Derivative of function with respect to other function, Successive differentiation, Differentiation using first principles.
15. **Limits and Derivatives:** Derivative introduced as rate of change both as that of distance function and geometrically, Intuitive idea of limit, Fundamental theorems on limits, Limits of polynomials and rational functions trigonometric, exponential and logarithmic functions, Definition of derivative relate it to slope of tangent of the curve, Derivative of sum, difference, product and quotient of functions, Derivatives of polynomial and trigonometric functions, Derivative of composite functions, Parametric functions, Derivative of function with respect to other function, Successive differentiation, Differentiation using first principles.
16. **Continuity, Differentiability and Application of Derivatives:** Continuity and differentiability, derivative of composite functions, chain rule, derivatives of inverse trigonometric functions, derivative of implicit functions. Concept of exponential and logarithmic functions.
 - Derivatives of logarithmic and exponential functions. Logarithmic differentiation, derivative of functions expressed in parametric forms. Second order derivatives. Rolle's and Lagrange's Mean Value Theorems (without proof) and their geometric interpretation.
 - Applications of derivatives: rate of change of bodies, increasing/decreasing functions, tangents and normals, use of derivatives in approximation, maxima and minima (first derivative test motivated geometrically and second derivative test given as a provable tool). Simple problems (that illustrate basic principles and understanding of the subject as well as real-life situations).
17. **Determinants:** Determinant of a square matrix (up to 3×3 matrices), properties of determinants, minors, cofactors and applications of determinants in finding the area of a triangle. Adjoint and inverse of a square matrix. Consistency, inconsistency and number of solutions of system of linear equations by examples, solving system of linear equations in

two or three variables (having unique solution) using inverse of a matrix.

18. **Matrices:** Concept, notation, order, equality, types of matrices, zero and identity matrix, transpose of a matrix, symmetric and skew symmetric matrices, Conjugate of matrix, Orthogonal matrix, idempotent matrix and Involutory matrix, Operation on matrices : Addition and multiplication and multiplication with a scalar. Simple properties of addition, multiplication and scalar multiplication. Non-commutativity of multiplication of matrices and existence of non-zero matrices whose product is the zero matrix (restrict to square matrices of order 2 and 3). Concept of elementary row and column operations. Invertible matrices and proof of the uniqueness of inverse, if it exists; (Here all matrices will have real entries).
19. **Integral and its Applications:** Integration as inverse process of differentiation. Integration of a variety of functions by substitution, by partial fractions and by parts, Evaluation of simple integrals of the different types and problems based on them. Definite integrals as a limit of a sum, fundamental Theorem of Calculus (without proof). Basic properties of definite integrals and evaluation of definite integrals. Applications in finding the area under simple curves, especially lines, circles/ parabolas/ellipses (in standard form only), Area between any of the two above said curves (the region should be clearly identifiable).
20. **Differential Equations:** Definition, order and degree, general and particular solutions of a differential equation. Formation of differential equation whose general solution is given. Solution of differential equations by method of separation of variables, solutions of homogeneous differential equations of first order and first degree. Solutions of linear differential equation of the type. $\frac{dy}{dx} + py = q$, where p and q are functions of x or constants. $\frac{dx}{dy} + px = q$, where p and q are functions of y or constants.
21. **Mathematical Reasoning:** Mathematically acceptable statements, Connecting words/phrases - consolidating the understanding of "if and only if (necessary and sufficient) condition", "implies", "and/or", "implied by", "and", "or", "there exists" and their use through variety of examples related to real life and Mathematics, Tautology, Contradiction and Duality, Algebra of statements, Validating the statements involving the connecting words, difference among contradiction, converse and contrapositive
22. **Statistics:** Measures of Dispersion : Range & Mean deviation, Variance and standard deviation of ungrouped / grouped data, The Median, Quartiles, Decile, Percentile, Mode of grouped and ungrouped data, analysis of frequency distributions with equal means but different variances. Correlation, types of correlation, Covariance, Karl Pearson Coefficient of correlation, Regression and its analysis.
23. **Commercial Mathematics:** Annuity, Partnership, Bill of Exchange, Foreign Exchange.
24. **Probability:** Random experiments, Outcomes, Sample spaces (set representation), Events, Occurrence of events: 'Not', 'And' and 'Or' events, Exhaustive events and mutually exclusive events, Axiomatic (set theoretic) probability, Connections with other theories of earlier classes, Probability of an event, Probability of 'net', 'and' 'or' events. Conditional probability, Multiplication theorem on probability, Independent events, Total probability, Bayes' theorem, Random variable and its probability distribution, Mean and variance of random variable, Repeated independent (Bernoulli) trials and Binomial distribution, Laws of probability addition theorem, De Morgan's Law, Poisson's Distribution, Normal Distribution.




MATHEMATICS MNEMONICS

Sets, Relations and Functions

Sets And Representations (a)

Today's Scenario, Equally Talented Singers Find Infinite New Songs To Sing.



Interpretation :

Types of Sets :

1. **Empty or Null Set** - A set which has no element.
2. **Finite Set** - A set having finite number of elements.
3. **Infinite Set** - A set having infinite number of elements.
4. **Equivalent Set** - Two finite sets A and B are said to be equivalent if $n(A)=n(B)$.
5. **Equal Set** - Two sets A and B are equal if every element of A is in B.
6. **Singleton Set** - A sets having one element is called singleton set.

Sets And Representations (b)

Laws of Algebra of Statements :

Iacd and Icai are friends

Interpretation :

1. **Idempotent Law** -
 - (i) $(A \wedge A) \Leftrightarrow A$
 - (ii) $(A \vee A) \Leftrightarrow A$
2. **Associative Law** -
 - (i) $(A \wedge B) \wedge C \Leftrightarrow A \wedge (B \wedge C)$
 - (ii) $(A \vee B) \vee C \Leftrightarrow A \vee (B \vee C)$
3. **Commutative Law** -
 - (i) $A \vee B \Leftrightarrow B \vee A$
 - (ii) $A \wedge B \Leftrightarrow B \wedge A$
4. **Distributive Law** -
 - (i) $A \vee (B \wedge C) \Leftrightarrow (A \vee B) \wedge (A \vee C)$
 - (ii) $A \wedge (B \vee C) \Leftrightarrow (A \wedge B) \vee (A \wedge C)$
5. **Identity Laws** -
 - (i) $A \vee T \Leftrightarrow A$
 - (ii) $A \wedge F \Leftrightarrow F$

$$(iii) \quad A \vee T \Leftrightarrow T$$

$$(iv) \quad A \vee F \Leftrightarrow A$$

6. Complement Laws -

$$(i) \quad A \vee (\sim A) \Leftrightarrow T$$

$$(ii) \quad A \wedge (\sim A) \Leftrightarrow F$$

$$(iii) \quad \sim T \Leftrightarrow F$$

$$(iv) \quad \sim F \Leftrightarrow T$$

7. Absorption Law -

$$(i) \quad A \vee (A \wedge B) \Leftrightarrow A$$

$$(ii) \quad A \wedge (A \vee B) \Leftrightarrow A$$

$$(iii) \quad \sim(A \wedge B) \Leftrightarrow (\sim A) \vee (\sim B)$$


8. Involution Law -

$$(i) \quad \sim(\sim A) \Leftrightarrow A$$

Complex Numbers and Quadratic Equations




Iodine Equipment Shows Result Negative One



Interpretation: Complex numbers are expressed in the form of $a+ib$ where 'i' is an imaginary number called 'iota' and the value of iota is $\sqrt{-1}$

Types of Linear Inequalities



Interpretation :

1. **Numerical Inequality** - $3 < 5, 8 > 4$
2. **Literal or Variable Inequalities** - $x < 5, y > 8$
3. **Double Inequality** - $5 < x < 9, 3 < y < 10$

4. Strict Inequality- $x < 9, 5 < 10$
5. Slack Inequality- $x \geq 7, y \leq 9$
6. linear Inequality in One Variable- $x < 9, y > 12$
7. linear Inequality in Two Variable- $5x + 7y < 12$
8. Quadratic Inequality- $x^2 + 5x \leq 10$

Matrices and Determinants



Identity Matrix-

$$A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad \begin{matrix} a_{ij} = 0 \text{ when } i \neq j \\ a_{ij} = 1 \text{ when } i = j \end{matrix}$$



Zero Matrix-

$$A = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$




Singular Matrix

A square matrix is said to be singular matrix if determinant of matrix denoted by $|A|$ is zero otherwise it is non zero matrix



Inverse Of a Matrix



Interpretation : Singular & Non Singular Matrix -

if $|A| = 0$, then A is singular. Otherwise A is non-singular


Inverse of a Matrix -

Inverse of a Matrix exists if A is non- singular i.e. $|A| \neq 0$, and is given by

$$A^{-1} = \frac{1}{|A|} \text{adj } A$$



Properties Of $|A|$



Interpretation : Properties of $|A|$ -

- (i) $|A|$ remains unchanged, if the rows and columns of A are interchanged i.e. $|A| = |A'|$
- (ii) If any two rows (or columns) of A are interchanged, then the sign of $|A|$ changes.
- (iii) If any two rows (or Columns) of A are identical then $|A| = 0$

Principle of Mathematical Induction



San Francis **P**rincipal **OM** Invited
Parents

SFPOMIP

Principle of Mathematical Induction (B)

Provided **T**est **P**aper of **1st** Term

PTP(1)T

Principle of Mathematical Induction (C)

Also **T**est **P**aper of **Kth** Term

ATP(K)T

Principle of Mathematical Induction (D)

Then **T**est **P**aper of **(K+1)th** Term

TPTP(K+1)T

Principle of Mathematical Induction (E)

Hence **P**aper of **nth** is **T**rustworthy
For **A**ll **N**ecessary **N**umbers

HP(n)TFANN

Principle of Mathematical Induction (F)

SFPOMIP-Steps for Principle of
Mathematical Induction Proof

Interpretation :

Step1: Let $P(n)$ be a result or
statement formulated in terms of n
in a given equation.

Principle of Mathematical Induction (G)

PTP(1)T-Prove that $P(1)$ is true.

Interpretation :

Step2: Prove that $P(1)$ is true.

Principle of Mathematical Induction (H)

ATP(K)T-Assume that $P(K)$ is true.

Interpretation :

Step3: Assume that $P(k)$ is true.

Principle of Mathematical Induction (I)

TPTP(K+1)T-prove that $P(k+1)$ is
true.

Interpretation :

Step4: Using step 3, prove that
 $P(k+1)$ is true.

Principle of Mathematical Induction (J)

HP(n)TFANN - Hence, by the
principle of mathematical induction,
 $P(n)$ is true for all natural numbers n

Interpretation :


Step5: Thus, $P(1)$ is true and $P(k+1)$
is true whenever $P(k)$ is true. Hence,
by the principle of mathematical
induction, $P(n)$ is true for all natural
numbers n .

Sequence and Series



Relationship between **AM**, **GM** and
HM

Area **O**f **H**ouse in **S**quare **G**igameter



Arithmetic Progression (AP)




(a) N^{th} Term of Arithmetic Progression -

NOAP



ETAPN1BD




Limits Continuity and Differentiability



**L' Hospital's Rule for Evaluating
Limits**

Numerator fights with **D**enominator,
both are **C**ritical $\left(\frac{0}{0}, \frac{\infty}{\infty}\right)$, **Lao** Hospital,
Bulao Dr.



Interpretation :

if $\lim_{x \rightarrow a} \frac{f(x)}{g(x)}$ takes $\frac{0}{0}$ or $\frac{\infty}{\infty}$ form

$$\text{then } \lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

$$\text{where } f'(x) = \frac{df(x)}{dx} \text{ and } g'(x) = \frac{dg(x)}{dx}$$



Sandwich Theorem for Evaluating Limits

Likhil always uses **S**ame size **(L)** Middle bread to make **T**hree layer Sandwich

(Three functions $f(x)$, $g(x)$ & $h(x)$)

If $f(x) \leq g(x) \leq h(x)$ and


$$\lim_{x \rightarrow a} f(x) = L = \lim_{x \rightarrow a} h(x) \text{ Then } \lim_{x \rightarrow a} g(x) = L$$

Interpretation :



If $f(x) \leq g(x) \leq h(x) \forall x \in (\alpha, \beta) - \{a\}$

$$\text{and } \lim_{x \rightarrow a} f(x) = L = \lim_{x \rightarrow a} h(x) \text{ then } \lim_{x \rightarrow a} g(x) = L$$

where $a \in (\alpha, \beta)$



Mean Value Theorem & Rolle's Theorem



Interpretation :

Mean Value Theorem -

if $f: [a, b] \rightarrow \mathbb{R}$ Continuous on $[a, b]$ and differential on (a, b) , then \exists some c in (a, b) such that-


$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

Rolle's Theorem -

If $f: [a, b] \rightarrow \mathbb{R}$ continuous on $[a, b]$ and differentiable on (a, b) and $f(a) = f(b)$ then \exists some c in (a, b) s.t. $f'(c) = 0$




SeCond FundAmental Theorem of Definite Integration



fcc is small fashionable Clothes

Counter **F**or **a**dorable **d**resses




Interpretation :

Let f be a continuous function defined on a closed interval $[a, b]$ and F be an anti derivative of f . Then $\int_a^b f(x) dx = [F(x)]_a^b = F(b) - F(a)$, where a and b are called limit of Integration.

Differential Equations




Linear Differential Equations



SOLDE-YIF-EIQ-IFC

Son Of Liladhar Dixit Eklavya (SOLDE)



S — Solution

D — Differential

O — Of

E — Equation

L — Linear

Interpretation :


Differential equation is of the form $\frac{dy}{dx} + py = Q$,

where P and Q are constants or the function of 'x' is called a first order linear differential equations. Its solution is given as

$$Y.IF = \int Q.IF + C$$



Homogeneous Differential Equation



Interpretation :

Differential equation can be expressed in the


$$\text{form } \frac{dy}{dx} = f(x, y) \text{ or } \frac{dx}{dy} = g(x, y)$$

where $f(x,y)$ and $g(x,y)$ are homogeneous functions of sum is called a homogeneous Differential equation. These equations can be solved by substituting $y=vx$ so that dependent variable y is changed to another variable v , where v is some unknown function.

Coordinate Geometry



Equation of Straight Line in Various forms :




Interpretation :

- (1) Point Slope form :- $y - y_1 = m(x - x_1)$
- (2) Slope intercept form :- $y = mx + c$
- (3) Two point form :- $y - y_1 = \frac{y_2 - y_1}{x_2 - x_1}(x - x_1)$
- (4) Intercept form:- $\frac{x}{a} + \frac{y}{b} = 1$
- (5) Normal / Perpendicular form :- $x \cos \alpha + y \sin \alpha = P$
- (6) General Form :- $ax + bx + c = 0$



Area of Triangle




ArEa Of Triangle

Area Equal to $\frac{1}{2}$ (one) (two)

$$\text{Area} = \frac{1}{2} [x_1(y_2 - y_3) + x_2(y_3 - y_1) + x_3(y_1 - y_2)]$$

Eccentricity of conic Sections

English alphabet Counting Sequence


**Interpretation :**

Eccentricity of Conic Sections-


- (a) If $e=1$, the conic is called parabola.
- (b) If $0 < e < 1$, the conic is called ellipse.
- (c) If $e > 1$, the conic is called hyperbola.
- (d) If $e=0$, the conic is called circle.

Three Dimensional Geometry**Direction Cosines**

Dance Choreographer Prefer Dieting



1 glass L e M o N juice

**Direction Ratios**


Director Remo a Professional Dancer




Choreographer created



3 Lifetime Movies with New faces a b c

**Interpretation :**

Direction cosines of a line are the cosines of the angles made by the line with the positive directions of the co-ordinate axes. If l, m, n are the D. cs of a line, then $l^2 + m^2 + n^2 = 1$


Vector Algebra**Types Of Vectors (A)****Interpretation :****Types of Vectors-**

1. **Zero Vector** - Initial and terminal points coincide
2. **Unit Vector** - Magnitude is unity
3. **Coinitial Vectors** - Same initial points
4. **Collinear vectors** - Parallel to the same Line
5. **Equal Vectors** - Same magnitude and direction
6. **Negative of a vector** - Same magnitude, opp. direction


**Properties Of Vectors(B)**

"Neither choose East nor choose north, always choose North-East and save your time".

North

**Interpretation :**


The vector sum of two coinitial vectors is given by the diagonal of the parallelogram whose adjacent sides are given vectors.



$$\vec{AB} + \vec{AC} = \vec{AD}$$

**Properties Of Vectors(C)**

Aao Bnaye Circle Tirahe par

**Interpretation:**

The vector sum of the three sides of a triangle taken in order is $\vec{0}$ i.e.

$$\vec{AB} + \vec{BC} + \vec{CA} = \vec{0}$$

Statistics & Probability

Mutually Exclusive Events

MEE-Mutual Enemies Everywhere

Morning **E**vening **E**veryday **C**annot **O**ccur

↓ Mutually Exclusive Events cannot occur

Sametime

simultaneously

Interpretation :

Events A & B are called mutually exclusive events if occurrence of any one of them excludes occurrence of other event, i.e. they cannot occur simultaneously.

eg: A die is thrown. Event A=All even outcomes & events B=All odd outcomes. then, A & B are mutually exclusive events, they cannot occur simultaneously

Poisson Distribution

DPD – Directions for Pure Dishes

↓ Distribution-Poisson Distribution

Perfect Restaurant Established
near Lajpat Nagar In NCR Provides
Rich Quality Naan Rice
Equal
$$P(r) = \lim_{n \rightarrow \infty} {}^nC_r p^r q^{(n-r)}$$

Here LemoN Quinoa Is Costliest Pure Dish

Here $\lambda = nq$ is called Poisson Distribution

Normal Distribution

DND — Do Not Disturb


↓ Distribution Normal Distribution

Parking x Entry Ticket is One rs.

Per Single Root for 2 Persons and


Entry Point is Mid Half Zone 2.

Power Minus $\frac{1}{2}$
$$p(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2}\frac{x^2}{\sigma^2}}$$



Variance and standard deviation for ungrouped data-

(a) Standard deviation for ungrouped data-



Variance for ungrouped data

"Vedic Fundamentals Under Graduates

(Variance) (for) (Ungrouped data)

lagao Square me Distinction

(Square) (Standard) (Deviation)

number Paao"

$$\text{Variance} = (\text{Standard deviation})^2$$

Interpretation :

Standard deviation of ungrouped data :

S.D. of ungrouped data is the square root of squared deviation from the mean of data. It is denoted by the symbol " σ "

Variance for ungrouped data :

Variance for ungrouped data is defined as the square of S.D. It is denoted by " σ^2 "

Sum and Difference of two Angles



Shape And Design Of Three Alphabets

Sum and Difference of two Angles



Remember the shape of **s**
 $\sin(A \pm B) =$
 Same Sign as **inside** the bracket in the expansion

Remember the curve of **c**
 Opposite the Curve of **c**
 $\cos(A \pm B) =$
 Opposite Sign in the expansion

Remember the shape
 $\tan(A \pm B) =$
 Twin Sign in (same) numerator

Interpretation :

$$* \sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$* \cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$* \tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

Standard General Solution of Trigonometric Ratios



S G S T R

Shabd ka Ginti Se Trigonometric Rishta

$$\sin \theta = 0 \Leftrightarrow \theta = n\pi$$

$\rho(\pi)$
 $P(\pi)$

$$\cos \theta = 0 \Leftrightarrow \theta = (2n+1)\frac{\pi}{2}$$

$\rho(\pi)$
t-(t for two-2 times 2)

$$\tan \theta = 0 \Leftrightarrow \theta = n\pi$$

$\rho(\pi)$
 $P(\pi)$

Interpretation :

The solution consisting of all possible solutions of a trigonometric equation is called its general solution.

$$* \sin \theta = 0 \Leftrightarrow \theta = n\pi$$


$$* \cos \theta = 0 \Leftrightarrow \theta = (2n+1)\frac{\pi}{2}$$

$$* \tan \theta = 0 \Leftrightarrow \theta = n\pi$$

Mathematical Reasoning



Algebra of statements -



AND

Mere Naam Me Mera Sign
 \wedge (And Sign)

Or
 Mere Naam Me Mera Sign
 \vee (or Sign)

CHAPTER : 1 SETS AND REPRESENTATIONS - PART -I

Let A and B be two sets. If every element of A is an element of B, then A is called a subset of B and written as $A \subset B$ or $B \supset A$ (read as 'A' is contained in 'B' or 'B contains A'). B is called superset of A.

Note:

1. Every set is a subset and superset of itself.
2. If A is not a subset of B, we write $A \not\subset B$.
3. The empty set is the subset of every set.
4. Power Set: If A is a set with $n(A) = m$, then no. of subset in power set $n[P(A)] = 2^m$
e.g. Let $A = \{3, 4\}$, then subsets of A are ϕ , $\{3\}$, $\{4\}$, $\{3, 4\}$. Here, $n(A) = 2$ and number of subsets of $A = 2^2 = 4$.

The number of elements in a finite set is represented by $n(A)$, known as cardinal number.
Eg.: $A = \{a, b, c, d, e\}$ Then, $n(A) = 5$

A set is a collection of well-defined distinguished objects. The sets are usually denoted by capital letters A, B, C etc., and the members or elements of the set are denoted by lower case letters a, b, c etc.,. If x is a member of the set A, we write $x \in A$ (read as 'x' belongs to A) and if x is not a member of set A, we write $x \notin A$ (read as 'x' doesn't belong to A). If x and y both belong to A, we write $x, y \in A$.
Some examples of sets are: A: odd numbers less than 10
N: the set of all natural numbers
V: the vowels in the English alphabets
R: the set of all real numbers.

In this form, we write a variable (say x) representing any member of the set followed by property satisfied by each member of the set.
e.g.: The set A of all prime number less than 10 in set builder form is written as
 $A = \{x \mid x \text{ is a prime number less than } 10\}$
The symbol " \mid " stands for the word "such that". Sometimes, we use symbol ":" in place of symbol " \mid ".

A set which has no element is called null set. It is denoted by symbol ϕ or $\{\}$.
e.g: Set of all real numbers whose square is -1 .
In set-builder form: $\{x : x \text{ is a real number whose square is } -1\}$
In roster form: $\{\}$ or ϕ


A set which has finite number of elements is called a finite set. Otherwise, it is called an infinite set.
e.g.: The set of all days in a week is a finite set whereas the set of all integers, denoted by $\{\dots, -2, -1, 0, 1, 2, \dots\}$ or $\{x \mid x \text{ is integers}\}$ is an infinite set.
An empty set ϕ which has no element is a finite set is called empty or void or null set.

Two sets A and B are set to be equal, written as $A = B$, if every element of A is in B and every element of B is in A.
e.g.: (i) $A = \{1, 2, 3, 4\}$ and $B = \{3, 1, 4, 2\}$, then $A = B$
(ii) $A = \{x : x - 5 = 0\}$ and $B = \{x : x \text{ is an integral positive root of the equation } x^2 - 2x - 15 = 0\}$
Then $A = B$

In this form, we first list all the members of the set within braces (curly brackets) and separate these by commas.
e.g: The set of all odd natural number less than 10 in this form is written as: $A = \{1, 3, 5, 7, 9\}$
In roster form, every element of the set is listed only once. The order in which the elements are listed is immaterial.
e.g: Each of the following sets denotes the same set
 $\{1, 2, 3\}$, $\{3, 2, 1\}$, $\{1, 3, 2\}$

A set having one element is called singleton set.
e.g.: (i) $\{0\}$ is a singleton set, whose only member is 0.
(ii) $A = \{x : 1 < x < 3, x \text{ is a natural number}\}$ is a singleton set which has only one member which is 2.

Two finite sets A and B are said to be equivalent, if $n(A) = n(B)$. Note: equal set are equivalent but equivalent sets need not to be equal.
e.g.: The sets $A = \{4, 5, 3, 2\}$ and $B = \{1, 6, 8, 9\}$ are equivalent, but are not equal.



CHAPTER : 1 SETS AND REPRESENTATIONS - PART -II

A Venn diagram is an illustration of the relationships between and among sets, groups of objects that share something common. These diagrams consist of rectangle and closed curves usually circles. E.g.:

In the given venn diagram $U = \{1, 2, 3, \dots, 10\}$ universe set of which $A = \{2, 4, 6, 8, 10\}$ and $B = \{4, 6\}$ are subsets and also $B \subset A$

- For any set A, we have
(a) $A \cup A = A$, (b) $A \cap A = A$, (c) $A \cup \phi = A$, (d) $A \cap \phi = \phi$, (e) $A \cup U = U$
(f) $A \cap U = A$, (g) $A - \phi = A$, (h) $A - A = \phi$
- For any two sets A and B we have
(a) $A \cup B = B \cup A$, (b) $A \cap B = B \cap A$, (c) $A - B \subseteq A$, (d) $B - A \subseteq B$
- For any three sets A, B and C, we have
(a) $A \cup (B \cap C) = (A \cup B) \cap C$, (b) $A \cap (B \cup C) = (A \cap B) \cup C$
(c) $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$, (d) $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$
(e) $A - (B \cup C) = (A - B) \cap (A - C)$, (f) $A - (B \cap C) = (A - B) \cup (A - C)$

The union of two sets A and B, written as $A \cup B$ (read as A union B) is the set of all elements which are either in A or in B or in both. Thus, $A \cup B = \{x : x \in A \text{ or } x \in B\}$ clearly, $x \in A \cup B \Rightarrow x \in A$ or $x \in B$ and $x \notin A \cup B \Rightarrow x \notin A$ and $x \notin B$ e.g. If $A = \{a, b, c, d\}$ and $B = \{c, d, e, f\}$ then $A \cup B = \{a, b, c, d, e, f\}$

The intersection of two sets A and B, written as $A \cap B$ (read as 'A' intersection 'B') is the set consisting of all the common elements of A and B.

Thus, $A \cap B = \{x : x \in A \text{ and } x \in B\}$
Clearly, $x \in A \cap B \Rightarrow \{x \in A \text{ and } x \in B\}$ and $x \notin A \cap B \Rightarrow \{x \notin A \text{ or } x \notin B\}$.
e.g. If $A = \{a, b, c, d\}$ and $B = \{c, d, e, f\}$ Then $A \cap B = \{c, d\}$

Two sets A and B are said to be disjoint, if $A \cap B = \phi$ i.e. A and B have no common element. e.g. if $A = \{1, 3, 5\}$ and $B = \{2, 4, 6\}$ Then, $A \cap B = \phi$, so A and B are disjoint.

The set containing all objects of element and of which all other sets are subsets is known as **universal sets** and denoted by U.
e.g. : For the set of all integers, the universal set can be the set of rational numbers or the set R of real numbers

Venn Diagram
Algebra of sets
Universal Set

Sets and Representations

If A and B are two sets, then their difference $A - B$ is defined as:
 $A - B = \{x : x \in A \text{ and } x \notin B\}$
Similarly, $B - A = \{x : x \in B \text{ and } x \notin A\}$
e.g. If $A = \{1, 2, 3, 4, 5\}$ and $B = \{1, 3, 5, 7, 9\}$ then
 $A - B = \{2, 4\}$ and $B - A = \{7, 9\}$

The symmetric difference of two sets A and B, denoted by $A \Delta B$, is defined as $(A \Delta B) = (A - B) \cup (B - A)$
e.g. If $A = \{1, 2, 3, 4, 5\}$ and $B = \{1, 3, 5, 7, 9\}$ then
 $(A \Delta B) = (A - B) \cup (B - A) = \{2, 4\} \cup \{7, 9\} = \{2, 4, 7, 9\}$

- The set of natural numbers $N = \{1, 2, 3, 4, 5, \dots\}$
 - The set of integers $Z = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$
 - The set of irrational numbers, $T = \{x : x \in R \text{ and } x \notin Q\}$
 - The set of rational number $Q = \{x : x = \frac{p}{q}, p, q \in Z \text{ and } q \neq 0\}$
- Relation among these subsets are $N \subset Z \subset Q, Q \subset R, T \subset R, N \not\subset T$

Interval Notation

Let a and b be real numbers with $a < b$

Set of Real Numbers

- $\{x | a < x < b\}$
- $\{x | a \leq x < b\}$
- $\{x | a < x \leq b\}$
- $\{x | a \leq x \leq b\}$
- $\{x | x < a\}$
- $\{x | x \leq a\}$
- $\{x | x > a\}$
- $\{x | x \geq a\}$
- R

Interval Notation

- (a, b)
- $[a, b)$
- $(a, b]$
- $[a, b]$
- $(-\infty, b)$
- $[-\infty, b]$
- (a, ∞)
- $[a, \infty)$

Region on the real number line

If U is a universal set and A is a subset of U, then complement of A is the set which contains those elements of U, which are not present in A and is denoted by A' or A^c . Thus, $A^c = \{x : x \in U \text{ and } x \notin A\}$
e.g.: If $U = \{1, 2, 3, 4, \dots\}$ and $A = \{2, 4, 6, 8, \dots\}$ then $A^c = \{1, 3, 5, 7, \dots\}$

Properties of complement

Complement law:

$$(i) A \cup A' = U \quad (ii) A \cap A' = \phi$$

De Morgan's Law:

$$(i) (A \cup B)' = A' \cap B' \quad (ii) (A \cap B)' = A' \cup B'$$

Double Complement law:

$$(A')' = A$$

Law of empty set and universal set

$$\phi' = U \text{ and } U' = \phi$$

CHAPTER

1

SETS AND REPRESENTATIONS

Chapter Objectives

Sets and their representations, Empty set, Finite and Infinite sets, Equal sets, Subsets of a set of real numbers especially intervals (with notations), Power set, Universal set, Venn diagrams, Union and Intersection of sets, Difference of sets, Complement of a set, Properties of Complement set, Algebraic properties of Union, Intersection and Complement of sets

STUDY MATERIAL

I. Concept Clarified

1. Sets and their notation :

A set is a collection of well-defined and well distinguished objects of our perception or thought which are distinct from each other.

➤ Notations

The sets are usually denoted by capital letters A, B, C , etc. and the members or elements of the set are denoted by lowercase letters a, b, c , etc.

If x is a member of the set A , we write $x \in A$ (read as 'x belongs to A') and if x is not a member of the set A , we write $x \notin A$ (read as 'x does not belong to A'). If x and y both belong to A , we write $x, y \in A$.

➤ Standard Notation

\mathbb{N} : A set of natural number.

$\mathbb{Q}^+ / \mathbb{Q}^-$: A set of all positive/negative rational number.

\mathbb{W} : A set of whole number.

\mathbb{R} : A set of real number.

\mathbb{Z} : A set of integers.

$\mathbb{R}^+ / \mathbb{R}^-$: A set of all positive/negative real number.

$\mathbb{Z}^+ / \mathbb{Z}^-$: A set of all positive/negative Integers.

\mathbb{C} : A set of complex number.

\mathbb{Q} : A set of rational number.

➤ Representation of a set

Sets are represented in the following two ways:

- Roster form or Tabular form
- Set Builder form or Rule Method

Roster Form or Tabular Form

In this method a set is described by listing elements, separated by commas and enclose then by curly brackets. For example, the set A of all odd natural numbers less than 10 in the Roster form is written as : $A = \{1, 3, 5, 7, 9\}$

Note

- In roster form, every element of the set is listed only once.
- The order in which the elements are listed is immaterial. For example, each of the following sets denotes the same set $\{1, 2, 3\}$, $\{3, 2, 1\}$, $\{1, 3, 2\}$.

Set-Builder Form or Rule Method

In this case we write down a property or rule which gives us all the elements of the set by that rule.

For example, the set A of all prime numbers less than 10 in the set-builder form is written as $A = \{x \mid x \text{ is a prime number less than } 10\}$. The symbol ' \mid ' stands for the words 'such that'. Sometimes, we use the symbol ':' in place of the symbol ' \mid '.

For more details,
scan the code



For more details,
scan the code



➤ **Types of Sets**

1. **Null set or Empty set** : A set having no element is called an empty set or a null set or void set. It is denoted by ϕ or $\{\}$ e.g. a set $A = \{x : x \text{ is an integer whose square root is negative natural number}\}$

2. **Finite Set** : A set which has only finite number of elements is called a finite set.

Example : $A = \{x : x^2 + 5x + 6 = 0\}$

3. **Infinite set** : A set which has an infinite number of elements is called an Infinite set.

Example : $A = \{y = 2x : x \text{ and } y \in \mathbb{R}\}$ since x can be any real number so the set contains any real number. Since the number of elements in the set is not defined so the given set is an infinite set.

4. **Singleton set** : A set consisting of a single element is called a singleton set.

Example : set $A = \{3\}$. Collection of SUN in solar system.

5. **Order of a finite set** : The number of elements in a finite set is called the order of the set A and is denoted by $O(A)$ or $n(A)$. It is also called cardinal number of the set.

Example : The order of set $B = \{-2, -3\}$ is $O(B)$ or $n(B) = 2$.

6. **Equal sets** : Two sets A and B are said to be equal if every element of A is a member of B , and every element of B is a member of A . We write $A = B$, if sets A and B are equal and $A \neq B$ when A and B are not equal.

Example : Consider set $A = \{x : x^2 + 5x + 6 = 0\}$ and set $B = \{-2, -3\}$. Since, A and B both have exactly same elements, hence set A is equal to set B .

7. **Equivalent set** : Two finite sets A and B are equivalent if their number of elements are same i.e. $n(A) = n(B)$.

Example : Let set A contain the vowel in the English alphabet and set B is defined as $B = \{x^2 : 1 \leq x \leq 5, x \in \mathbb{N}\}$.
Since, $n(A) = n(B) = 5$, so they are equivalent set.

Note : Equal sets will always be equivalent but equivalent sets may not be equal sets.

8. **Subsets** : Let A and B be two sets. If every element of A is the element of B , then A is called a subset of B . If A is a subset of B , we write $A \subseteq B$.

9. **Proper subset** : If A is a subset of B and $A \neq B$ then A is a proper subset of B and we write $A \subset B$.

In other words, if A is a proper subset of B , then all elements of A are in B but B contains at least one element that is not in A .

Example : Let set $A = \{x^2 : 1 < x < 5, x \in \mathbb{N}\}$ and $B = \{x^2 : 2 \leq x \leq 4, x \in \mathbb{N}\}$, so $A \subseteq B$.

Example : Let set $A = \{x^2 : 1 < x < 5, x \in \mathbb{N}\}$ and $B = \{x^2 : 1 < x < 10, x \in \mathbb{N}\}$, so $A \subset B$

Note : Every set is a subset of itself i.e. $A \subseteq A$ for all A . Empty set is a subset of every set

$$\mathbb{N} \subset \mathbb{W} \subset \mathbb{Z} \subset \mathbb{Q} \subset \mathbb{R} \subset \mathbb{C}$$

The total number of subsets of a finite set containing n elements is 2^n .

A set is a subset of itself but not a proper subset of itself.

If A has n elements, then its power set $P(A)$ contains exactly 2^n elements.

10. **Power set** : Let A be any set. The set of all the subsets of A is called power set of A and is denoted by $P(A)$.

Example : Let set $A = \{a, b, c\}$, then Power Set of A is given as $P(A) = \{\{\}, \{a\}, \{b\}, \{c\}, \{a, b\}, \{a, c\}, \{b, c\}, \{a, b, c\}\}$

11. **Universal set** : A set consisting of all possible elements which occur in the fixed domain, is called a Universal set and is denoted by U . All sets are contained in the Universal set.

Example : Set of Real numbers is a universal set for natural numbers, whole numbers and rational numbers.

12. **Comparable sets** : Two sets A and B are comparable, if $A \subseteq B$ or $B \subseteq A$.

Example : Let two sets $A = \{x^2 : 1 < x < 5, x \in \mathbb{N}\}$ and $B = \{4, 9, 16\}$

Since, $A \subseteq B$ or $B \subseteq A$ so we can compare two sets.

13. **Venn diagram** : The diagram drawn to represent sets are called venn – diagrams, where universal set U is represented by rectangle and its subsets represented by closed curves within the rectangle.

➤ **Operations on Sets with Venn Diagram**

1. **Union of Two Sets** : The union of two sets A and B , written as $A \cup B$ (read as 'A union B'), is the set consisting of all the elements which are either in A or in B or in both.


For more details,
scan the code



For more details,
scan the code




Example : Let two sets $A = \{x^2 : 1 < x < 5, x \in \mathbb{N}\}$ and $B = \{1, 2, 3, 5\}$ then $A \cup B = \{1, 2, 3, 4, 5, 9, 16\}$



Combination of the shaded regions together represent the union of sets.

2. Intersection of Two sets : The intersection of two sets A and B , written as $A \cap B$ (read as 'A' intersection 'B') is the set consisting of all the common elements of A and B .

Example : Let two sets $A = \{x^2 : 1 < x < 5, x \in \mathbb{N}\}$ and $B = \{1, 2, 3, 4, 5\}$ then $A \cap B = \{4\}$.




The shaded region which is common to both the shaded regions represents intersection of sets

3. Disjoint Sets : Two sets A and B are called disjoint, if $A \cap B = \phi$. They do not have any common element.

Example : Let two sets $A = \{x^2 : 1 < x < 5, x \in \mathbb{N}\}$ and $B = \{1, 5, 6\}$


Since, there are no elements in common, therefore, $A \cap B = \phi$



4. Difference of Two Sets : If A and B are two sets, then their difference $A - B$ is defined as $A - B = \{x : x \in A \text{ and } x \notin B\}$. Similarly, $B - A = \{x : x \in B \text{ and } x \notin A\}$.

Example : Let two sets $A = \{x^2 : 1 < x < 5, x \in \mathbb{N}\}$ and $B = \{1, 2, 3\}$ Then, $A - B = \{4, 9, 16\}$

5. Complement of a set : Let A is a subset of universal set U , then the complement of A with respect to U is the set of all those element of U which are not in A . It is denoted by A' or A^c or $U - A$. The union of a set A and its complement A' gives the universal set U of which A and A' are a subset, i.e., $A \cup A' = U$



Also, the intersection of a set A and its complement A' gives the empty set ϕ . i.e., $A \cap A' = \phi$

Law of Double Complementation : According to this law if we take the complement of the complemented set A' then, we get the set A itself. i.e., $(A')' = A$

Law of empty set and universal set : According to this law the complement of the universal set gives the empty set and vice-versa i.e. $U' = \phi$ and $\phi' = U$.

➤ LAWS OF ALGEBRA OF SETS

For three sets A , B and C

1. Idempotent Law

- (a) $A \cup A = A$
- (b) $A \cap A = A$

2. Identity Law

- (a) $A \cup \phi = A$
- (b) $A \cap U = A$

3. Commutative Law

- (a) $A \cup B = B \cup A$
- (b) $A \cap B = B \cap A$

4. Associative Law

- (a) $(A \cup B) \cup C = A \cup (B \cup C)$
- (b) $A \cap (B \cap C) = (A \cap B) \cap C$

5. Distributive Law

- (a) $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$
- (b) $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$

6. De-Morgan's Law

- (a) $(A \cup B)' = A' \cap B'$

For more details,
scan the code



$$(b) (A \cap B)' = A' \cup B'$$

7. Symmetric difference property

$$A \Delta B = (A - B) \cup (B - A)$$

8. Results on operation on sets

- (a) $A - (B \cap C) = (A - B) \cup (A - C)$
- (b) $A - (B \cup C) = (A - B) \cap (A - C)$
- (c) $A - B = A \cap B'$
- (d) $B - A = B \cap A'$
- (e) $A - B = A \Leftrightarrow A \cap B = \phi$
- (f) $(A - B) \cup B = A \cup B$
- (g) $(A - B) \cap B = \phi$
- (h) $A \cap B \subseteq A$ and $A \cap B \subseteq B$
- (i) $A \cup (A \cap B) = A$
- (j) $A \cap (A \cup B) = A$

9. More on symmetric difference property

$$(a) (A - B) \cup (B - A) = (A \cup B) - (A \cap B)$$

For more details,
scan the code



(b) $A \cap (B - C) = (A \cap B) - (A \cap C)$

(c) $A \cap (B \Delta C) = (A \cap B) \Delta (A \cap C)$

(d) $(A \cap B) \cup (A - B) = A$

(e) $A \cup (B - A) = A \cup B$

10. Universal Set property

(a) $U' = \phi$

(b) $\phi' = U$

(c) $(A')' = A$

(d) $A \cap A' = \phi$

(e) $A \cup A' = U$

(f) $A \subseteq B \Leftrightarrow B' \subseteq A'$

II. Important Formulae

(1) $n(A \cup B) = n(A) + n(B) - n(A \cap B)$

(2) $n(A \cup B) = n(A) + n(B) \Leftrightarrow A, B$ are disjoint non – void sets.

(3) $n(A - B) = n(A) - n(A \cap B)$

(4) $n(B - A) = n(B) - n(A \cap B)$

(5) $n(A \Delta B) = n(A) + n(B) - 2n(A \cap B)$

(6) $n(A \cup B \cup C) = n(A) + n(B) + n(C) - n(A \cap B) - n(B \cap C) - n(A \cap C) + n(A \cap B \cap C)$

(7) number of elements in exactly two of the sets A, B, C $= n(A \cap B) + n(B \cap C) + n(C \cap A) - 3n(A \cap B \cap C)$

(8) number of elements in exactly one of the sets A, B, C

$$= n(A) + n(B) + n(C) - 2n(A \cap B) - 2n(B \cap C) - 2n(A \cap C) + 3n(A \cap B \cap C)$$

(9) $n(A' \cup B') = n((A \cap B)') = n(U) - n(A \cap B)$


(10) $n(A' \cap B') = n((A \cup B)') = n(U) - n(A \cup B)$

For more details,
scan the code


CHAPTER - 2 : RELATIONS AND FUNCTIONS - PART- I

The composition of functions $f : A \rightarrow B$ and $g : B \rightarrow C$ is denoted by $g \circ f$, and is defined as $g \circ f : A \rightarrow C$ given by $g \circ f(x) = g(f(x)) \forall x \in A$. e.g. let $A = N$ and $f, g : N \rightarrow N$ such that $f(x) = x^2$ and $g(x) = x^3 \forall x \in N$. Then $g \circ f(2) = g(f(2)) = g(2^2) = 4^3 = 64$.

A function $f : X \rightarrow Y$ is invertible, if \exists a function $g : Y \rightarrow X$ such that $g \circ f = I_X$ and $f \circ g = I_Y$. Then, g is the inverse of f . If f is invertible, then it is both one-one and onto and vice-versa.
eg. If $f(x) = x$ and $f : N \rightarrow N$, then f is invertible.
Theorem 1 : If $f : X \rightarrow Y, g : Y \rightarrow Z$ and $H : Z \rightarrow S$ are functions, then $h \circ (g \circ f) = (h \circ g) \circ f$.
Theorem 2 : Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be two invertible functions, then $g \circ f$ is invertible with $(g \circ f)^{-1} = f^{-1} \circ g^{-1}$.



CHAPTER : 2 RELATIONS AND FUNCTIONS - PART- II



CHAPTER

2

RELATIONS AND FUNCTIONS

Chapter Objectives

Ordered pairs, Cartesian product of sets, Number of elements in the Cartesian product of two finite sets, Cartesian product of the set of real with itself (up to $\mathbb{R} \times \mathbb{R} \times \mathbb{R}$), Definition of relation, pictorial diagrams, domain, co-domain and range of a relation, types of relation, binary operation, Types of relations : reflexive, symmetric, transitive and equivalence relations. One to one and onto functions, composite functions, inverse of a function. Binary operations.

Function as a special type of relation, Pictorial representation of a function, domain, co-domain and range of a function, real valued functions, domain and range of these functions, constant, identity, polynomial, rational, modulus, signum, exponential, logarithmic and greatest integer functions, with their graphs, composition of function, sum, difference, product and quotients of functions.

STUDY MATERIAL

I. Concept Clarified

1. Ordered Pair :

An ordered pair consists of two objects or elements in a given fixed order. If $a \in A$ and $b \in B$, then all pairs in the form (a, b) is called the ordered pair.

Example : Let $2^a = 7^b$. the solution of this set that is $(0, 0)$ will be ordered pair where $a = 0$ and $b = 0$.

Equality of ordered pairs : Two ordered pairs (a_1, b_1) and (a_2, b_2) are equal, if $a_1 = a_2$ and $b_1 = b_2$.

2. Cartesian product of sets

For two sets A and B (non-empty sets), the set of all ordered pairs (a, b) such that $a \in A$ and $b \in B$ is called cartesian product of the sets A and B, denoted by $A \times B$.

$A \times B = \{(a, b) : a \in A \text{ and } b \in B\}$

Example : Let Set $A = \left\{ f(x)f(4) - \frac{1}{2} \left(f\left(\frac{x}{4}\right) + f(4x) \right) \text{ where } f(x) = \cos \log x \right\}$ and

Set $B = \{b, c \text{ where } f(x) = bx^2 + cx + d \text{ and } f(x+1) - f(x) = 8x + 3\}$

The element Sets are,

$$\text{Set } A = \left\{ \begin{aligned} &f(x)f(4) - \frac{1}{2} \left(f\left(\frac{x}{4}\right) + f(4x) \right) \text{ where } f(x) = \cos \log x \\ &\Rightarrow \cos(\log x) \cos(\log 4) - \frac{1}{2} \left(\cos \log \frac{x}{4} + \cos \log 4x \right) \\ &\Rightarrow \cos(\log x) \cos(\log 4) - \frac{1}{2} [2 \cos(\log x) \cos(\log 4)] \\ &\Rightarrow 0 \end{aligned} \right\}$$

$$\text{Set } A = \{0\}$$

$$\text{Set } B = \left\{ \begin{aligned} &b, c \text{ where } f(x) = bx^2 + cx + d \text{ and } f(x+1) - f(x) = 8x + 3 \\ &b = 4, c = -1 \end{aligned} \right\}$$

$$\text{Set } B = \{4, -1\}$$

Cartesian product of $A \times B$ is $\{(0, 4)(0, -1)\}$ where $0 \in \text{Set } A$ and $4, -1 \in \text{Set } B$.

For more details,
scan the code



3. Number of elements in the Cartesian product of two finite sets :

$$n(A \times B) = n(A) \cdot n(B)$$

4. Cartesian product of the set of real with itself ($R \times R \times R$)

The set $R \times R \times R$ represent the coordinates of all the points in three dimensional plane.

$$\text{Set } B = \left\{ \begin{array}{l} b, c \text{ where } f(x) = bx^2 + cx + d \text{ and } f(x+1) - f(x) = 8x + 3 \\ b = 4, c = -1 \end{array} \right\}$$

$$\text{Set } B = \{4, -1\}$$


Then Cartesian product of $B \times B \times B$ will contain $\{(4, 4, 4), (4, 4, -1), (4, -1, 4), (-1, 4, 4), (4, -1, -1), (-1, 4, -1), (-1, -1, 4), (-1, -1, -1)\}$

These are also known as ordered triplet.

5. Relation

If A and B are two non-empty sets then any subset of $A \times B$ is called relation from A to B. Such relations between two non-empty sets is called binary relation and if R is a relation from a to b and $(a, b) \in R$, then it is written as $a R b$ and read as a is related to b.


Example : Consider a set $A = \{x : x^3 - 6x^2 + 11x - 6 = 0\}$ and set B contains elements as $\text{Set } B = \{x : x^3 - 12x^2 + 44x - 48 = 0\}$. Then the relation between Set A and B from A to B will be set of any combinations from set A to set B.

**6. Domain And Range of A Relation :**

Let R be a relation from a set A to set B. Then, set of all first components or x-coordinates of the ordered pairs belonging to R is called the domain of R, while the set of all second components or y-coordinates of the ordered pairs belonging to R is called the range of R.

Thus, domain of $R = \{a : (a, b) \in R\}$ and range of $R = \{b : (a, b) \in R\}$

Example $R = \{y = x^2 - 3, \text{ where } x \in \mathbb{N} < 4\}$, then the domain of the function is $\{1, 2, 3\}$ correspondingly the range of the function is $\{-2, 1, 6\}$. The graph of the following relation is as follow.



For more details,
scan the code

**7. Types of Relation****(i) Void Relation :**

As $\phi \subset A \times A$, for any set A, so ϕ is a relation on A, called the empty or void relation.

Example : The relation R on the set $A = \{1, 2, 3, 4\}$ defined by $R = \{(a, b) : a + b = 10\}$ since $a + b \neq 10$ for any two elements of set A. Therefore, $(a, b) \notin R$ for any $a, b \in A \Rightarrow R$ does not contain any element $A \times A$

$\Rightarrow R$ is an empty set.

$\Rightarrow R$ is the void relation on A.

(ii) Universal Relation :

Since, $A \times A \subseteq A \times A$, so $A \times A$ is a relation on A, called the universal relation.

Example : Relation on the set $A = \{1, 2, 3, 4, 5, 6\}$ by $R = \{(a, b) \in R : |a - b| \geq 0\}$ since $|a - b| \geq 0$ for all $a, b \in A$

$\Rightarrow (a, b) \in R$ for all $(a, b) \in A \times A$.

\Rightarrow each element of set A is related to every element of set A .

$\Rightarrow R = A \times A$.

$\Rightarrow R$ is a universal relation on set A .

(iii) Identity Relation :

The relation $I_A = \{(a, a) : a \in A\}$ is called the identity relation on A .

Example : suppose $A = \{1, 2, 3\}$, then the set of ordered pairs $\{(1, 1), (2, 2), (3, 3)\}$ is the identity relation on set ' A '.

(iv) Reflexive Relation :

A relation R is said to be reflexive relation, if every element of A is related to itself. Thus,

$(a, a) \in R, \forall a \in A \Rightarrow R$ is reflexive.

Example : A relation R is defined on the set Z (set of all integers) by " $a R b$ if and only if $2a + 3b$ is divisible by 5", for all $a, b \in Z$. So let say, $a \in Z$. Now $2a + 3a = 5a$, which is divisible by 5. Therefore $a R a$ holds for all a in Z i.e. R is reflexive.

(v) Symmetric Relation :

A relation R is said to be symmetric relation, iff

$(a, b) \in R \Rightarrow (b, a) \in R, \forall a, b \in A$

i.e. $a R b \Rightarrow b R a, \forall a, b \in A$

$\Rightarrow R$ is symmetric. The set A of natural numbers.

Example : A relation R is defined on the set Z by " $a R b$ if $a - b$ is divisible by 5" for $a, b \in Z$. So let say $a, b \in Z$ and $a R b$ hold. Then $a - b$ is divisible by 5 and therefore $b - a$ is divisible by 5.

Thus, $a R b \Rightarrow b R a$ and therefore R is symmetric.

(vi) Transitive Relation :

A relation R is said to be transitive relation, iff $(a, b) \in R$ and $(b, c) \in R$

$\Rightarrow (a, c) \in R, \forall a, b, c \in A$

Example : in the set A of natural numbers if the relation R be defined by ' x less than y ' then $a < b$ and $b < c$ imply $a < c$, that is, $a R b$ and $b R c \Rightarrow a R c$. Hence this relation is transitive.

(vii) Equivalence Relation :

A relation R is said to be an equivalence relation, iff it is simultaneously reflexive, symmetric and transitive on A .

Example : Let assume that R be a relation on the set R of real numbers defined by $x R y$ if and only if $x - y$ is an integer. Prove that R is an equivalence relation on R .

Reflexive : Consider x belongs to R , then $x - x = 0$ which is an integer. Therefore $x R x$.

Symmetric : Consider x and y belongs to R and $x R y$. Then $x - y$ is an integer. Thus, $y - x = -(x - y)$, $y - x$ is also an integer. Therefore $y R x$.

Transitive : Consider x and y belongs to R , $x R y$ and $y R z$. Therefore $x - y$ and $y - z$ are integers. According to the transitive property, $(x - y) + (y - z) = x - z$ is also an integer. So that $x R z$.

Thus, R is an equivalence relation on R .

(viii) Partial Order Relation :

A relation R is said to be a partial order relation, iff it is simultaneously reflexive, anti symmetric and transitive on A .

Example : $(4, 2) \in R$ and $(2, 1) \in R$, implies $(4, 1) \in R$. As the relation is reflexive, anti-symmetric and transitive. Hence, it is a partial order relation.

(ix) Equivalence classes of an equivalent relation.

Let R be equivalence relation in A ($\neq \phi$). Let $a \in A$. Then, the equivalence class of a denoted by $[a]$ or

$\{\bar{a}\}$ is defined as the set of all those points of A which are related to a under the relation R .

Example : $f(x)$ is the set of all integers, we can define the equivalence relation \sim by saying ' $a \sim b$ if and only if $(a - b)$ is divisible by 9'. Then the equivalence class of 4 would include $-32, -23, -14, -5, 4, 13, 22$, and 31 .

For more details,
scan the code



For more details,
scan the code



For more details,
scan the code



For more details,
scan the code



For more details,
scan the code



For more details,
scan the code



(x) Composition of relation :

Let R and S be two relations from sets A to B and B to C respectively, then we can define relation composition of S over R or SoR from A to C such that $(a, c) \in SoR \Leftrightarrow \exists b \in B$ such that $(a, b) \in R$ and $(b, c) \in S$.

This relation $S \circ R$ is called the composition of R and S .


(a) $RoS \neq SoR$

(b) $(SoR)^{-1} = R^{-1} \circ S^{-1}$ known as reversal rule.

Example : Let $X = \{4, 5, 6\}$, $Y = \{a, b, c\}$ and $Z = \{l, m, n\}$. Consider the relation R_1 from X to Y and R_2 from Y to Z .

$$R_1 = \{(4, a), (4, b), (5, c), (6, a), (6, c)\}$$

$$R_2 = \{(a, l), (a, n), (b, l), (b, m), (c, l), (c, m), (c, n)\}$$



The composition of relation (i) $R_2 \circ R_1$

The composition relation $R_2 \circ R_1$ as shown in fig :




Fig : $R_1 \circ R_2$

$$R_2 \circ R_1 = \{(4, l), (4, n), (5, l), (5, m), (5, n), (6, l), (6, m), (6, n)\}$$

(xi) Congruence Modulo m :

Let m be an arbitrary but fixed integer. Two integers a and b are said to be congruence modulo m , if $a - b$ is divisible by m and we write $a \equiv b \pmod{m}$.

Example : $a \equiv b \pmod{m} \Leftrightarrow a - b$ is divisible by m . For example $26 \equiv 11 \pmod{5} \Leftrightarrow 26 - 11$ is divisible by 5.

For more details,
scan the code




For more details,
scan the code

**8. Binary Operation :**

Let S be a non-empty set. A function f from $S \times S$ to S is called a binary operation on S i.e. $f : S \times S \rightarrow S$ is a binary operation on set S .

Example : Consider the binary operation \wedge on the set $\{1, 2, 3, 4, 5\}$ defined by $a \wedge b = \min \{a, b\}$.


For more details,
scan the code



$\begin{smallmatrix} a \\ b \end{smallmatrix}$	1	2	3	4	5
1	1	1	1	1	1
2	1	2	2	2	2
3	1	2	3	3	3
4	1	2	3	4	4
5	1	2	3	4	5

9. Function :


A function is like a machine which gives unique output for each input that is fed into it. But every machine is designed for certain defined inputs for e.g. a washing machine is designed for washing cloths and not the wood. Similarly the functions are defined for certain inputs which are called as its **domain** and corresponding outputs are called **Range**.



Let A and B be two sets and let there exist a rule or manner or correspondence ' f ' which associates to each element of A to a unique element in B, then f is called a **Function** or **Mapping** from A to B. It is denoted by symbol

$$f:(A,B) \text{ or } f:A \rightarrow B \text{ or } A \xrightarrow{f} B$$

Which reads ' f is a function from A to B' or ' f maps A to B'.



Note : Every function is a relation but every relation is not necessarily a function.

10. Domain and Range of function :

For a relation from set A to set B i.e. aRb , **all the elements of set A are called as the domain of the relation R while all the elements of set B are called as the co-domain of the relation R.**

Range is the set of all second elements from the ordered pairs (a, b) in the relation aRb .

$$\text{Domain of } f = \{a \mid a \in A, (a, f(a)) \in f\}$$

$$\text{Range of } f = \{f(a) \mid a \in A, f(a) \in B, (a, f(a)) \in f\}$$

For the relation aRb , domain is considered as the input to relation R while the co-domain is the possible outputs and range is the actual output.

Example : Domain (D) and Range (R) of $f(x) = \sin^{-1}(x)$ is $[-1, 1]$ and $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ respectively.

11. Types of function :

(a) Polynomial Function : A function f is defined by

$$f(x) = a_n x^n + a_{n-1} x^{n-1} + a_{n-2} x^{n-2} + \dots + ax + a_0$$

For more details,
scan the code





where n is a non negative integer and $a_n, a_{n-1}, \dots, a, a_0$ are real number and $a_n \neq 0$, then f is called a polynomial function of degree n . A polynomial function is always continuous.

(b) Rational Function :

A function of type, $f(x) = \frac{g(x)}{h(x)}$, where $g(x)$ and $h(x)$ are polynomials functions and $h(x) \neq 0$, is called rational function.

(c) Exponential Function :

A function $f(x) = a^x$ where ($a > 0, a \neq 1, x \in \mathbb{R}$) is called an exponential function. $f(x) = a^x$ is called an exponential function because the variable x is the exponent. It should not be confused with power function $g(x) = x^2$ in which variable x is the base. For $f(x) = e^x$ domain is \mathbb{R} and range is \mathbb{R}^+ .





If $a > 1$ then f is increasing function.

If $0 < a < 1$ then f is strictly decreasing function.

If $a \neq 0$ then f is strictly monotonic function

(d) Logarithmic Function :

If $a > 0$ and $a \neq 1$ then a function of the form $f(x) = \log_a x, x > 0$ is called general logarithmic function $D_f = (0, \infty)$.




(e) Absolute Value Function (or Modulus Function) :

A function $y = f(x) = |x|$ is called the absolute value function or modulus function. It is defined as :

$$y = |x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$$

For $f(x) = |x|$, domain is \mathbb{R} and Range is $\mathbb{R}^+ \cup \{0\}$. See below for its figure.



(f) Greatest Integer Or Step up Function :

The function $y = f(x) = [x]$ is called the greatest integer function where $[x]$ denotes the greatest integer less than or equal to x . Note that for :

For more details,
scan the code

